
BOVAX® 3V





AGRIPROM PAKISTAN (Pvt) Ltd.

Plot No.19-B, OFF Abdul Sattar Eidhi Road, Near Qazalbash Chowk, Lahore-Pakistan Tel: +92 42 35232901 +92 300 080 1249 info@agriprompakistan.com.pk www.agriprompakistan.com.pk



BOVAX° 3V

Trivalent inactivated vaccine against respiratory diseases in cattle due to the IBR and Pl₃ viruses and Pasteurella multocida

Efficacy

- ✓ It induces high and prolonged antibody levels.
- ✓ It antagonises the primary agents of respiratory infections (viruses) and secondary agents (the bacteria responsible for superinfections), amongst which the most dangerous is *Pasteurella multocida*.
- ✓ The adjuvant used is a highly purified saponin, which reduces local reactions to a minimum and possesses high immunostimulant properties.

Safety

BOVAX® 3V IS AN INACTIVATED VACCINE:

- ✓ It does not cause persistent infection due to the vaccine strain of IBR virus, as happens when animals immunized with live attenuated vaccines.
- ✓ There is no risk of spread of the vaccine strains at the farm.
- ✓ May be used in animals of any age or production category.
- ✓ Does not cause decreases in fertility or productivity, side eff ects which can accompany immunisation performed with attenuated strains.
- Does not cause vaccination stress, consequently eliminating the risk of secondary infections.





In case vaccination is introduced in the farm for the first time, inject all the subjects at the same time, with a booster after 2-4 weeks, then follow the programmes recommended.

If the animals are moved to a different farm, it is advisable to perform basic vaccination programme, namely the first two vaccinations or at least the first vaccination, prior to transport.

خصوصیات سانس کی تین مہلک بیاریوں گل گھوٹو، پیراافلو ئنزا، آئی بی آر کے خلاف قوت مدافعت پیدا کرتی ہے۔ استعمال خوراک: 5ml فی جانور زیر جلدانحیک کریں اورانحیکشن لگانے سے پہلے بو تل کواچھی طرح ہلائیں۔ و سیسینیشن پر و گرام گوشت والا جانور: ۔ پہلاانحیکشن فارم پر پہنچنے پر دو سراانحیکشن پہلے انحیکشن سے 2-4 ہفتے بعد اور ہر 6 ماہ بعد بوسٹر ڈوز لگوائیں۔ دوردھ دینے والے جانور: ۔ (کٹاکٹی / وچھاوچھی) پہلاانحیکشن پیدا ہونے کے تین ہفتے بعد

و یکسین کو فرتج میں دو سے آٹھ ڈگری سینٹی گریڈتک رکھیں اور ہر گزبرف بننے سے بچائیں ویکسین کو سورج کی روشن سے محفوظ رکھیں۔

COMPOSITION

Each 5 ml dose of vaccine contains: inactivated culture of infectious bovine rhinotracheitis (IBR) virus: 10⁷ DICT₅₀, inactivated culture of parainfluenza 3 virus: 10⁷ DICT₅₀, inactivated culture of *Pasteurella multocida*: 15x10⁹ CFU. Adjuvant: alhydrogel saponin (Quil A).

INDICATIONS

Active immunization against infectious bovine rhinotracheitis (IBR), parainfluenza 3 and pasteurellosis in cattle.

ADMINISTRATION

Subcutaneous.

DOSAGE

The vaccine should be administered by subcutaneous injection at a dosage of 5 ml.

In farms where treatment is being initiated treat all animals at the same time.

Administer booster 2-4 weeks later.

Beef cattle:

- first vaccination on arrival, followed by a booster injection 2-4 weeks later;
- second booster after about 6 months.

Dairy and breeding cattle: Calves:

- first vaccination at 3rd week of life, followed by a booster 2-4 weeks later; further boosters every 6-8 months.

Heifers and cows:

vaccinations every 6-8 months.



Produced according to Good Manufacturing Practices of European Union (EU-GMP), the best certified quality system for veterinary biologicals

پیکنگ: 250 ml



Plot No.19-B, OFF Abdul Sattar Eidhi Road, Near Qazalbash Chowk, Lahore-Pakistan Tel: +92 42 35232901 +92 300 080 1249 info@agriprompakistan.com.pk www.agriprompakistan.com.pk

